# **REVIEW**

**Open Access** 

# Retinopathy of prematurity: from oxygen management to molecular manipulation



Jonathan Woods<sup>1\*</sup> and Susmito Biswas<sup>2</sup>

# Abstract

**Introduction** Retinopathy of prematurity (ROP) is a vasoproliferative disorder of the premature retina with the potential to progress to extraretinal neovascularisation. This review serves as an introduction to retinopathy of prematurity (ROP), outlining key parts of ROP pathophysiology, diagnosis and treatment. ROP is traditionally diagnosed by indirect ophthalmoscopy and classified using anatomical zones, stages of disease, and the presence or absence of "plus disease" (dilation and tortuosity of the major retinal arterioles and venules). ROP has a bi-phasic pathophysiology: initial hyperoxia causes reduced retinal vascularisation, followed by pathological vaso-proliferation resulting from subsequent hypoxia and driven by vascular endothelial growth factor (VEGF).

**Advancements in management** This review summarises previous trials to establish optimum oxygen exposure levels in newborns and more recently the development of anti-VEGF agents locally delivered to block pathological neovascularisation, which is technically easier to administer and less destructive than laser treatment.

**Future directions** There remains an ongoing concern regarding the potential unwanted systemic effects of intravitreally administered anti-VEGF on the overall development of the premature baby. Ongoing dosing studies may lessen these fears by identifying the minimally effective dose required to block extraretinal neovascularisation.

Keywords Infant, Newborn, Retinopathy of prematurity, Ophthalmoscopy, Cell proliferation, Oxygen

# Introduction

Preterm birth has a potentially wide-ranging impact on neurovascular development, including the neonatal retina. The incidence of premature birth is increasing globally and varies internationally (5–18% in some European and African states, respectively) [1]. Preterm survival is also increasing globally; this progress is more marked in countries with rapidly improving neonatal facilities [2] that are sufficiently developed to increase preterm survival, but insufficiently developed to reduce associated morbidity [3]. There is, therefore, an increasing number

\*Correspondence:

<sup>2</sup> Manchester Royal Eve Hospital, Manchester University Hospital NHS

of preterm babies at risk of lifelong disability due to complications associated with prematurity.

Retinopathy of prematurity (ROP) is a vasoproliferative disorder of the premature retina with the potential to progress to extraretinal neovascularisation. This can subsequently lead to retinal detachment and sight loss. A recent multicentre audit in the UK revealed that 4% of babies with a birth weight (BW) < 1500 g required treatment for ROP [4]. The incidence of ROP in settings with less developed neonatal facilities is higher, since ROP can develop in babies of greater birth weight and gestational age (GA) in these settings [3, 5].

# Prematurity and normal eye development Retinal vascular development

The retina consists of organised layers of photoreceptors, interneurons, glial elements, and their various interconnections, overlying the retinal pigment epithelial layer. The retina receives oxygen through a dual blood supply with



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

Jonathan Woods

JAW911@student.bham.ac.uk

<sup>&</sup>lt;sup>1</sup> University of Birmingham Medical School, Medical School, College of Medical and Dental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

Foundation Trust, Oxford Rd, Manchester M13 9WL, UK

the outer layers receiving a supply from the choroidal layer and the inner retina predominantly receiving a supply from the retinal vasculature. Vascularisation of the retina starts in the second month of gestation from the optic disc and moves peripherally behind a "wave of neuronal differentiation" and consequent increasing metabolic activity [6], reaching the nasal side in the eighth month and the temporal side at approximately the time of term delivery.

# The role of oxygen in the pathogenesis of ROP

In utero, the partial pressure of oxygen  $(PO_2)$  in the umbilical vein is less than 50 mmHg [7]. Thus, retinal vasculature develops in a controlled, relatively hypoxic environment. Following delivery, arterial PO<sub>2</sub> rises; neonates given supplementary oxygen may be exposed to yet higher, more fluctuating levels of oxygen. Crosse and Jameson observed in 1952 that the incidence of ROP in neonates rose and fell with the liberal and restricted administration of oxygen, respectively [8]. The ELGAN study corroborates that such hyperoxia on blood gases contributes to the development of ROP [9]. Blood transfusion containing adult haemoglobin may further increase oxygen availability in developing structures due to the different oxygen affinities of foetal and adult haemoglobin [10]. The preterm retina is particularly vulnerable to hyperoxia for two reasons: the immature vasculature has a reduced ability to autoregulate [11], and the reserve of antioxidants is not large enough to protect from damage by reactive oxygen species [12] due to prevention of full transfer of  $\varpi$ -long-chain polyunsaturated fatty acids (LCPUFAs) from mother to baby in the third trimester [13]. Limited data from clinical trials indicate the potential for supplementation of LCPUFAs in the early post-natal period may reduce the incidence of severe ROP [14-16].

# Levels of insulin-like growth factor-1 (IGF-1) in the premature neonate

Reduced concentrations of IGF-1, seen in preterm neonates [17], is a key contributor to ROP development due to its synergistic action with VEGF where angiogenesis occurs only when IGF-1 reaches a critical level [18]. The deficit in IGF-1 means that vitreoretinal VEGF levels continue to build until IGF-1 reaches the critical level to support angiogenesis. The supranormal levels of VEGF then result in extraretinal neovascularisation [19]. The possibility that supplemental IGF-1 given from 24 h of birth until reaching 30 weeks postmenstrual age (PMA) might prevent severe ROP was tested in a clinical trial with intravenous infusions of supplemental recombinant human IGF-1 combined with recombinant human IGF-1 binding protein. Unfortunately, no reduction in severe ROP was observed in the treatment arm compared to the control arm although there was a reduced incidence of bronchopulmonary dysplasia in the treatment arm. Further studies are required to clarify the potential of IGF-1 [20].

#### **Risk factors**

Understanding retinal development in the context of prematurity provides an explanation for the risk factors associated with ROP: gestational age, birth weight, postnatal weight gain and nutrition linked to IGF-1 [21] and oxygen exposure [9]. Therefore, ROP may feature alongside other conditions associated with disrupted vascular development or abnormal fluctuations in oxygen saturation such as poor brain growth, bronchopulmonary dysplasia and necrotising enterocolitis.

Dysregulation of blood vessel development due to infection or inflammation, such as chorioamnionitis, may also form a "prephase" of ROP [22]. It is unknown whether intra-uterine events that may cause preterm delivery predispose to the development of ROP. Higher incidence in males and Caucasian babies suggests there may also be a genetic predisposition as implied by ethnic variations in incidence of severe ROP [23].

# Pathogenesis

Retinopathy of prematurity can be thought of in two phases: the cessation of normal vascular development in high-oxygen environments, followed by pathological compensatory vascularisation of the retina [24].

#### Phase one (hyperoxia)

In addition to decreased levels of IGF-1 and maternal  $\omega$ -LCPUFAs, exposure of the premature retina to increased oxygen causes further dysregulation of vascular development by suppression of hypoxia-induced factors (HIFs) which reduces levels of angiogenic factors such as vascular endothelial growth factor (VEGF) and erythropoietin (EPO) [25]. In extreme cases, an oxygeninduced retinopathy has been demonstrated in pre-term babies ( $\geq$ 28 weeks and  $\geq$  1000 g) given unmonitored and unblended 100% oxygen for 3 days or longer after birth [26].

#### Phase two (hypoxia)

As the retina develops, its metabolic demand increases such that the environment becomes hypoxic. In hypoxic conditions, the breakdown of HIFs is reduced [27, 28]. HIFs upregulate the expression of VEGF which acts via specific receptors (VEGFR-1 and VEGFR-2) [29] to cause increased migration, proliferation and permeability of endothelial cells. In ROP, this process becomes pathological due to the excessive amounts of VEGF in the absence of sufficient IGF-1 to support angiogenesis. Once IGF-1 reaches sufficient levels, the excess VEGF causes extraretinal neovascularisation which subsequently undergoes cicatrisation. This cicatricial process is accompanied by contracture of fibrous metaplastic tissue which, if extensive enough, can lead to a tractional retinal detachment [18, 30, 31]. Whilst the role of EPO in ROP is not well elucidated, excessive concentrations of EPO in the vitreous have also been correlated with levels of vitreous VEGF and partial retinal detachment [32].

#### Timing of phase transition

The transition from phase 1 to phase 2 has a significant bearing on the timing of initiation for screening of ROP, the purpose of which is to identify sight-threatening ROP before it becomes untreatable. Whilst this is not precisely known in any given premature baby, we know from natural history studies that there is a close association of ROP development to post-menstrual age [33]. In babies born before 27 weeks of gestation, the earliest onset of stage 3 ROP is not earlier than 31 weeks, and therefore, this can be used as a guide to initiate screening in the most premature neonates [34]. Figure 1 shows a summary diagram of the progression of ROP (Fig. 1).

# **Diagnosis and classification**

ROP is traditionally diagnosed by indirect ophthalmoscopy but, more recently, is increasingly being identified through digital retinal imaging [35]. ROP is classified using the International Classification of ROP (ICROP3) standards [36] which denote ROP in terms of zones, stages, clock hours (extent) and "plus disease". The retina is divided into three concentric zones (Fig. 2). The stages are defined by the appearance of the retinal vasculature at the junction of the vascularised retina with the non-vascularised retina. Initially, this may appear as just incomplete vascularisation (immature vascularisation), previously termed stage 0. Stage 1 represents a pale demarcation line at the junction between vascularised and avascular retina, stage 2 represents an elevation of the demarcation line (ridge) and stage 3 represents the development of extraretinal neovascularisation. Small angioblastic buds, termed "popcorn", may appear above the stage 2 ridge presaging the development of stage 3. Changes in the appearance of the blood vessels at the posterior pole within zone I may be noted during the progression of ROP with gradually increasing vessel dilation and tortuosity of the major arterioles and venules, termed "plus disease". This is a significant driver for treatment for ROP and is therefore a critical feature. There is a gradation in the progression of vessel tortuosity and dilation from the absence of plus disease to plus disease with intermediate levels being termed "pre-plus". Distinguishing plus disease from pre-plus is often a matter of judgement and there is considerable subjectivity and, consequently, interobserver variation, but is likely to be aided in the future with the use of artificial intelligence algorithms [37]. However, it is possible that the apparent size and shape of vessels become distorted by the pressure change associated with using contact wideangle retinal imaging camera lenses [38]. Stage 3 ROP in the presence of plus disease forms a fibrovascular complex that may progress to a tractional retinal detachment.



Fig. 1 The 2-phase model of ROP. Diagrams shown below the timeline are cartoons of retinal flat mounts, similar to those seen in rat and mouse model



**Fig. 2** Zones of the right eye and appearance of retinal vasculature according to ROP stage. Zone I, the most posterior, is a circular zone centred on the optic nerve and with a radius twice the distance (2x) from the centre of the optic nerve to the centre of the macula (x). Zone II represents a middle zone, concentric to zone I, extending from the border of zone I to the nasal ora serrata (furthest extent of the nasal retina). Zone II is itself subdivided into a posterior zone II, two disc diameters out from the border of zone I, and anterior zone II encompassing the remaining peripheral area of zone II. Zone III is the most peripheral zone extending from the temporal border of zone II and capturing a remaining temporal crescentic area beyond zone II. ROP is likely to be more severe when located in zone I or posterior zone II. Stages 1–3 can occur in any zone of the eye. Here, they are shown in zone II for clarity

Where there is a partial detachment, this is termed stage 4: if not involving the macula, this is stage 4a and if involving the macula, this is stage 4b. Stage 5 denotes a total detachment of the retina and is further subdivided into stage 5a (total retinal detachment); stage 5b (total detachment of the retina with a closed funnel configuration); stage 5c (accompanying changes to the anterior segment of the eye such as cataract, glaucoma and corneal opacity). Stages 4 and 5 are typically associated with poor visual outcomes and blindness.

#### Implications for management

As ROP has two phases, there is a lag phase from birth to the development of the ROP lesion. A balance therefore needs to be struck between screening unnecessarily early and commencing in a timely way in order to identify preterm infants at risk of sight-threatening ROP. UK guidelines recommend that eye examinations are not carried out before 31 weeks PMA [39], due to observational studies showing that severe ROP is not seen before this point [34]. Accurate classification of ROP is not only important for identifying babies that require treatment, but is also for identifying infants that are at higher risk of progression and therefore requiring more frequent monitoring intervals. Table 1 shows how zone, stage and the presence of plus disease inform types of ROP. Treatment-warranted ROP is termed "type 1". "Type 2" ROP represents ROP status that requires careful monitoring, on a weekly basis. Babies that have immature vascularisation only in anterior zone II, stage 1 in anterior zone II or stage 2 in anterior zone II without any plus or pre-plus disease are all deemed to be lower risk and may be monitored at two weekly intervals. Immature vascularisation terminating in zone I or posterior zone II; or stage 1 or 2 ROP in zone I or posterior zone II or stage 3 ROP in Zone II (type 2 ROP); the presence of any pre-plus disease, all increase the risk of progression and would necessitate more frequent, weekly monitoring [39].

Aggressive ROP (A-ROP) is a rare but severe form of ROP. This typically develops in zone I or posterior zone II, but may be located in more anterior zones in some parts of the world with limited resources available for optimal neonatal care. The hallmark of A-ROP is the marked prominence of plus disease typically in all

Table 1	Using stage, zone	e, and presence	e of "plus disease"	to classify ROP i	into type 1	and type 2
---------	-------------------	-----------------	---------------------	-------------------	-------------	------------

	Zone I		Zone II		Zone III	
Stage 1	⊕ = <i>Type</i> 1	$\Theta = Type 2$	Examination interva disease status	ils based on plus /pre-plus	Examination intervals based on plus / pre-plus disease status	
Stage 2	$\oplus = Type$ 1	$\Theta = $ Type 2	⊕ = <i>Туре</i> 1	Examination intervals based on pre-plus disease status	Examination intervals based on plus / pre-plus disease status	
Stage 3	$\oplus = Type$ 1	$\Theta = Type \ 1$	⊕ = <i>Туре</i> 1	$\Theta =$ Type 2	Examination intervals based on plus / pre-plus disease status	

 $\oplus$  Plus disease

 $\ominus$  No plus disease

4 retinal quadrants of the eye with a poorly defined, flat neovascularisation not associated with a demarcation line or ridge.

#### Prevention and treatment

#### Oxygen management

Between 1994 and 2014, large, multi-centre randomised control trials (RCTs)—STOP-ROP [40], SUPPORT [41], BOOST II [42], and COT [43]—have been conducted across developed countries. Table 2 compares these studies (Table 2).

Collectively, these studies demonstrate the balance of supplementary oxygen: higher oxygen saturation targets are associated with ROP and bronchopulmonary dysplasia, whereas a lower target is associated with higher rates of neurological damage, necrotising enterocolitis and death, which caused early termination of the BOOST II trial. Currently, UK guidelines recommend a target oxygen saturation of 91–95% [44]. The 2-phase model of the pathogenesis of ROP may suggest that there is no single target SaO<sub>2</sub>. Rather, when considering ROP in isolation, target SaO<sub>2</sub> should ideally change over time as the disease moves from its hyperoxic to the hypoxic stage. However, this should be carefully balanced against the systemic implications of changing oxygen saturations and practical challenges of implementation.

#### Pharmacological approaches

Currently, there are two dominant modalities for the treatment of ROP, (1) intravitreal injection of anti-VEGF antibodies and/or (2) laser ablation of the avascular retina. Laser therapy requires prolonged sedation and, even when initiated early as in the Early Treatment of ROP (ETROP) study, cannot prevent retinal detachment in 12% of cases [45]. Thus, increasing focus has been given to a pharmacological approach which would have the benefit of bedside administration, reduced cost and would reduce the burden on specialist ophthalmologists experienced in delivering laser treatments [46].

The antibody bevacizumab binds to all forms of VEGF-A [47]. Intravitreal injection of bevacizumab has been widely used in other vasoproliferative ocular diseases such as age-related macular degeneration, diabetic retinopathy, central and branch retinal vein occlusion in adult patients [48–51]. In addition, intravitreal bevacizumab has been used in children for conditions such as Coats disease and subretinal neovascular membranes. However, its use in all ocular diseases in both adults and children is off-label. Study data shows that, in the UK, Bevacizumab is used by some ophthalmologists to treat A-ROP and posterior type-1 ROP [4].

BEAT-ROP, a multicentre RCT, compared the efficacy of bevacizumab against standard laser treatment, using recurrence of ROP before 54 weeks postmenstrual age as a primary outcome [52]. The authors concluded that the anti-VEGF agent was shown to be effective against stage 3+ disease in zone I, but not in zone II and larger numbers of participants were required to comment on safety. However, these conclusions have been questioned, since using recurrence at 54 weeks as a primary outcome may miss later reactivation of ROP giving a false impression of bevacizumab's efficacy [53]. The figure of 54 weeks was based on the recurrence of ROP following laser treatment for ROP from the ETROP study but does not take into account the possibility of late reactivation of ROP that is seen in babies treated with anti-VEGF (see below).

A Cochrane Review [54] of anti-VEGF agents for the treatment of ROP included four RCTs (including BEAT-ROP) comparing bevacizumab with laser therapy. The authors concluded that aggregated data suggest bevacizumab as monotherapy does not reduce recurrence or risk of retinal detachment (although, as suggested by BEAT-ROP, treatment for zone I ROP is promising).

Safety concerns have been raised by clinicians regarding the use of anti-VEGF in this age group [53, 55]. Studies in humans show that bevacizumab, when injected intra-ocularly, can escape into the general circulation [56]. Furthermore, primate models demonstrate that the serum concentration of bevacizumab remains raised for

	STOP-ROP	Support	Boost II	COT
Location	30 centers in USA	22 centers in USA	Centers in Australia ( $n$ = 9), New Zea- land ( $n$ = 5), and UK ( $n$ = 20)	25 centers in Canada, USA, Argentina, Finland, Germany, Israel
Recruitment period	1994–1999	2005-2009	2006–2010	2006-2010
Latest follow-up	3 months after due date	Discharge or 36 weeks (latest)	2 years	18 months
Inclusion criteria	Preterm infants AND confirmed ROP in $\geq 1$ eye AND median SaO <sub>2</sub> < 94%	GA: 24–26 weeks +6 days	GA < 28 weeks	GA: 23–27 weeks + 6 days
Number of participants	649	1316	2448	1201
Target oxygen saturations	89–94% 96–99%	85–89% 91–95%	85–89% 91–95%	85–89% 91–95%
Mortality	2.2% 2.8%	19.9% 16.2%	23.1% 15.9%	16.6% 15.3%
	No RR/OR available	RR= 1.27 (95% CI= 1.01-1.60)	RR = 1.45 (95% CI = 1.15-1.84)*	OR= 1.11 (95% CI = 0.80-1.54)
Development of severe ROP	48% 41%	8.6% 17.9%	10.6% 13.5%	12.8% 13.1%
	OR = 0.72 (95% CI = 0.52 - 1.01)	RR=0.52 (95% CI=0.37-0.73)	RR = 0.79 (95% CI = 0.63-1.00)	OR=0.95 (95% CI=0.65-1.39)
Other differences in outcomes	Pneumonia/Chronic lung disease 4.7% higher in higher target SaO <sub>2</sub> group	85–89% group had a decreased rate of $O_2$ use at 36 weeks GA ( $p$ = 0.002)	Risk of necrotising enterocolitis 2.4% higher in 85–89% group (RR = 1.31, 95% CI = 1.01–1.68)	No significant differences in brain injury, necrotising enterocolitis or sev- ered BPD
Conclusions	SaO <sub>2</sub> has no significant effect on risk of developing ROP SaO <sub>2</sub> > 96% may have pathological pulmonary effects	Significantly increased mortality among 85–89% group No significant difference in risk of ROP	Significantly increased risk of death in infants with target SaO <sub>2</sub> < 90% *Recruitment stopped early	No significant differences in mortality or disability (including ROP)

 Table 2
 Summary of studies comparing different target SaO2 for prevention of ROP

OR Odds ratio (adjusted), Cl Confidence interval, RR Relative risk (adjusted), BPD Bronchopulmonary dysplasia

up to 8 weeks [57]. The 2018 Cochrane review [54] of anti-VEFG agents concluded that insufficient safety data existed for the recommendation for routine use of bevacizumab. Roohipoor et al. [58] recruited 116 participants, randomised to 0.625 mg bevacizumab or laser therapy. The study found that ROP regressed in all cases of the bevacizumab arm (5 eyes—3.2%—requiring retreatment at a mean of 6.01 weeks after initial treatment) and 97.3% of eyes in the laser therapy arm (p=0.20). Despite follow-up to 90 weeks, only ocular complications were recorded in the bevacizumab arm (one instance of cataract formation).

The dose of bevacizumab used in Roohipoor's and the BEAT-ROP study was 0.625 mg, half of the adult dose. However, this may be significantly more than required to neutralise VEGF in the vitreous. Further studies looking at lower doses of Bevacizumab ranging from 0.16 to 0.004 mg have shown promising outcomes [59, 60], although there is a likelihood of reactivation occurring sooner with lower doses [61].

An alternative anti-VEGF antibody, Ranibizumab, a human monoclonal antibody fragment derived from the parent antibody Bevacizumab [47] has been licenced and approved for treatment of retinal vasoproliferative disorders in adults and also for ROP by the National Health Service (NHS England) [62] and European Medicines Agency (EMA) [63] following completion of the RAINBOW trial [64]-225 participants, randomized to ranibizumab 0.2 mg, ranibizumab 0.1 mg, and laser photocoagulation therapy. Analysis showed that ranibizumab 0.2 mg may be superior to laser therapy (treatment success odds ratio compared to laser therapy=2.19. 95% CI 0.99-4.82). Stahl et al. record that "Death, serious and non-serious systemic adverse events, and ocular adverse events were evenly distributed between the three groups" at 24 weeks [64]. One notable advantage that Ranibizumab may have over Bevacizumab is the reduced duration of systemic VEGF suppression [56, 65, 66].

Longer-term follow-up from the CARE-ROP study [67] which compared different doses of ranibizumab (0.2 mg and 0.12 mg) shows no significant difference in ophthalmic and neurodevelopmental outcomes between the treatment arms after 1 and 2 years, respectively.

A third agent, Aflibercept, has the potential to prevent VEGF-driven pathogenesis by binding VEGF-A, VEGF-B as a soluble decoy receptor [68]. Aflibercept is licensed for use in adult retinal vasoproliferative disorders by the US Food and Drug Administration FDA [69] and EMA [70]. The FIREFLEYE study tested Aflibercept in a multicentre, international non-inferiority RCT for treatment of type 1 ROP and A-ROP with an active control comparator group of babies treated with laser therapy [71]. As with the RAINBOW study, the primary outcome was the proportion of infants without active ROP and unfavorable structural outcomes (retinal detachment, macular dragging, macular fold or retrolental opacity) 24 weeks after starting treatment. One hundred eighteen infants were randomized, and 113 were treated in a 2:1 ratio of Aflbercept (0.4 mg) (75 babies) or laser (38 babies). Although treatment success was slightly higher in the Aflibercept-treated arm, the credible interval for the treatment difference was not greater than the prespecified value of -5% and therefore non-inferiority could not be proven. Further and larger studies of Aflibercept are therefore required to reach any definitive conclusions regarding the comparative effect of Aflibercept to laser in type 1 ROP.

ROP regression following anti-VEGF is a notable for its rapidity. The RAINBOW study documented plus disease regressing significantly faster, at a median time of 4 days (interquartile range (IQR) of 3-8 days) following 0.2 mg of Ranibizumab compared to a median time of 16 days (IQR of 9–22 days) following laser. Similarly, stage 3 was noted to regress at a median of 8 days (IQR 5-17 days) following 0.2 mg of Ranibizumab versus a median of 16 days (IQR 9-29 days) following laser. Indeed, A-ROP was noted to take nearly 3 times longer to regress following laser treatment versus Ranibizumab [72]. This study also showed that incomplete regression of ROP requiring retreatment was more frequently observed in lasertreated eyes with around 22% of eyes requiring further treatment at a median interval of 15 days post-laser compared to 8% of Ranibizumab-treated eyes at a median interval of 21 days after injection [72].

Significant concerns around treatment with anti-VEGF include the prolonged follow-up required to identify reactivation of the disease, that is, a recurrence of acute phase features of ROP, and the management of any persistent avascular retina in the periphery, which arises from interruption of vascular development following intravitreal anti-VEGF administration. Reactivation may occur after complete or incomplete regression of ROP after anti-VEGF injections. It rarely occurs after complete regression of ROP following laser treatment. Reactivation following anti-VEGF treatment may involve, at its mildest, the recurrence of stage 1 demarcation line or more severe features such as stage 3 and/or plus disease. The natural history of such reactivations may not follow through the typical stages of ROP either and may just show features of plus disease associated with fine neovascularisation [34]. Such reactivation has been described more than 12 months after treatment with anti-VEGF (Bevacizumab) [73]. In the RAINBOW study, 31% of infants receiving 0.2 mg of intravitreal Ranibizumab had one or more additional treatments during their follow-up with a median time to first retreatment of 55 days (range

29–111 days) [64]. Ranibizumab may be associated with more rapid reactivation than bevacizumab or aflibercept [74]. Reactivation of ROP following intravitreal anti-VEGF injection may also be seen more in more posterior aggressive diseases or in those injected at a younger post-menstrual age [75]. Very low-dose bevacizumab for treatment of ROP, ranging from 0.002 to 0.016 mg, is also associated with earlier reactivations with a mean of 76.4 days to reactivation compared to low-dose bevacizumab, ranging from 0.031 to 0.25 mg, with a mean of 85.7 days to reactivation [61].

Babies treated with anti-VEGF may never fully vascularise to the peripheral retina leaving a persistent area of avascular retina (PAR) that runs the risk of late reactivation of ROP. This potential for late reactivation in a persistent avascular zone leaves a number of management dilemmas. PAR occurs in untreated, spontaneously regressed ROP as well as in ROP treated with anti-VEGF. Long-term pathological changes within the PAR observed in adult patients with a history of ROP that never reached treatment criteria include atrophic retinal holes, retinal tears, both of which can lead to retinal detachment, tractional retinoschisis (splitting of the retinal layers), vascular anomalies including telangiectatic vessels, vascular leakage and neovascularisation [76]. Similarly fundus fluorescein angiography studies carried out in eyes treated with a history of anti-VEGF injections with PAR demonstrate vascular anomalies such as vascular leakage, shunts, abnormal vessel branching and tangles [77]. There is no consensus regarding the management of such cases in terms of treatment or followup regimes. Additionally, in the short to medium term, to continue to undertake regular, serial ROP examinations require parents or guardians to return with their baby for ongoing retinal examinations that may pose practical difficulties, particularly if they have to travel a significant distance. Furthermore, the examinations become increasingly stressful and difficult to carry out in infants with increasing age. Further research is required to answer the questions of whether and when to treat or not.

# Conclusion

ROP is a significant concern in preterm infants and progression to severe disease has a poor visual prognosis. Its pathogenesis can be understood in the context of premature transition from intra- to extra-uterine environments and consequent disruption to oxygen availability in the retina which dictates vascular development via signalling molecules. When considering ROP in isolation, reducing oxygen availability in the retina may help to control this process. However, the implications of this on other developing structures such as the brain and gastrointestinal tract and higher mortality are unacceptable.

Other significant research questions include refining the criteria for identifying babies at risk of serious ROP. The relationship of IGF-1, post-natal weight gain and severe ROP is well established [78] and has encouraged the development of a number of algorithms based on GA, BWT and weight gain and ethnicity [79, 80]. Implementation of refined and validated screening algorithms may reduce the overall burden of screening.

The increasing uptake of digital retinal imaging and subsequent capability of image analysis also lends itself to the development of artificial intelligence and deep learning algorithms to assist clinicians in identifying critical features of ROP, such as plus disease versus no plus or pre-plus disease, vascular severity scores or stage of ROP. Many are showing promise but as yet are not reliable enough to entirely replace the experienced clinician [81].

Interventions such as laser ablation of the avascular retina persist, but more recently anti-VEGF agents offer a more targeted approach. Future optimal dosage studies may eradicate long-term wider safety fears and elucidate optimum dosages or therapeutic agents specific to the stage and zone of ROP. Further studies may also address the problem of management of the peripheral avascular retina following anti-VEGF treatment. Going forward, alternative ways of targeting VEGF, such as the use of the beta-blocker, propranolol [82] and the targeting of other pathological pathways that lead to ROP, such as LCPUFAs and IGF-1 may warrant further investigation.

#### Abbreviations

A-ROP	Aggressive Retinopathy of Prematurity
BEAT-ROP	Bevacizimab Eliminates the Angiogenic Threat of Retinopathy of
	Prematurity
BOOST II	Benefits of Oxygen Saturation Targeting
BW	Birth weight
CARE-ROP	Comparing Alternative Ranibizumab Dosages for Safety and Effi-
	cacy in Retinopathy of Prematurity
COT	Canadian Oxygen Trial
ELGAN	Extremely low gestational age newborns
EMA	European Medicines Agency
ETROP	Early Treatment of Retinopathy of Prematurity
EPO	Erythropoietin
FDA	Food and Drug Administration (US)
FIREFLEYE	Aflibercept for Retinopathy of Prematurity—Intravitreal Injection
	Versus Laser Therapy (FIREFLEYE)
GA	Gestational age
HIFs	Hypoxia-induced factors
ICROP3	International Classification of Retinopathy of Prematurity 3
IGF-1	Insulin-like growth factor-1
LCPUFAs	Long-chain polyunsaturated fatty acids
NHS	National Health Service (UK)
PAR	Persistent avascular retina
PMA	Postmenstrual age
PO2	Partial pressure of oxygen

RAINBOW	RAnibizumab Compared with Laser Therapy for the Treatment of
	INfants BOrn Prematurely With Retinopathy of Prematurity
RCTs	Randomized control trials
ROP	Retinopathy of prematurity
SaO2	Oxygen saturation
STOP-ROP	Supplemental Therapeutic Oxygen for Prethreshold Retinopathy
	of Prematurity
SUPPORT	Surfactant, Positive Pressure, and Pulse Oximetry Randomized
	Trial
VEGF	Vascular endothelial growth factor
VEGFR	Vascular endothelial growth factor receptor

#### Acknowledgements

Dr. Robert Barry and Dr. Gibran Farook Butt provided advice during the background research of this review. Figures 1 and 2 created with BioRender.com.

#### Authors' contributions

 $\mathsf{JW}$  and  $\mathsf{SB}$  wrote and reviewed the manuscript.  $\mathsf{JW}$  prepared all figures and tables.

#### Funding

The authors declare that no funds, grants or other support were received during the preparation of this manuscript.

#### Availability of data and materials

This review used published literature, accessed through the University of Birmingham Library.

# Declarations

#### Ethics approval and consent to participate

This review of existing literature did not require ethical approval.

#### **Competing interests**

The authors declare no competing interests.

Received: 24 November 2022 Accepted: 11 August 2023 Published online: 15 September 2023

#### References

- Blencowe H, Cousens S, Oestergaard MZ, Chou D, Moller AB, Narwal R et al (2012) National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. Lancet 379(9832):2162–2172. https://doi.org/10.1016/S0140-6736(12)60820-4
- Liu L, Oza S, Hogan D, Chu Y, Perin J, Zhu J et al (2016) Global, regional, and national causes of under-5 mortality in 2000–15: an updated systematic analysis with implications for the sustainable development goals. Lancet 388(10063):3027–3035. https://doi.org/10.1016/S0140-6736(16) 31593-8
- Gilbert C, Fielder A, Gordillo L, Quinn G, Semiglia R, Visintin P et al (2005) Characteristics of infants with severe retinopathy of prematurity in countries with low, moderate, and high levels of development: implications for screening programs. Pediatrics 115(5):e518–e525. https://doi.org/10. 1542/peds.2004-1180
- Adams G, Bunce C, Xing W, Butler L, Long V, Reddy A et al (2017) Treatment trends for retinopathy of prematurity in the UK: active surveillance study of infants at risk. BMJ Open 7(3):e013366. https://doi.org/10.1136/ bmjopen-2016-013366
- Bowe T, Nyamai L, Ademola-Popoola D, Amphornphruet A, Anzures R, Cernichiaro-Espinosa LA et al (2019) The current state of retinopathy of prematurity in India, Kenya, Mexico, Nigeria, Philippines, Romania, Thailand, and Venezuela. Digit J Ophthalmol 25(4):49–58. https://doi.org/ 10.5693/djo.01.2019.08.002
- Schoenwolf GC, Bleyl SB, Brauer PR, Francis-West PH (2015) Larsen's human embryology, 5th edn. Churchill Livingstone, Philadelphia, p 554

- Nicolaides KH, Economides DL, Soothill PW (1989) Blood gases, pH, and lactate in appropriate- and small-for-gestational-age fetuses. Am J Obstet Gynecol 161(4):996–1001. https://doi.org/10.1016/0002-9378(89)90770-9
- Crosse VM, Evans PJ (1952) Prevention of retrolental fibroplasia. Arch Ophthalmol 48(1):83–87. https://doi.org/10.1001/archopht.1952.00920 010086012
- Hauspurg AK, Allred EN, Vanderveen DK, Chen M, Bednarek FJ, Cole C et al (2011) Blood gases and retinopathy of prematurity: the ELGAN study. Neonatology 99(2):104–111. https://doi.org/10.1159/000308454
- Wang YC, Chan OW, Chiang MC, Yang PH, Chu SM, Hsu JF et al (2017) Red blood cell transfusion and clinical outcomes in extremely low birth weight preterm infants. Pediatr Neonatol 58(3):216–222. https://doi.org/ 10.1016/j.pedneo.2016.03.009
- Milligan DW (1980) Failure of autoregulation and intraventricular haemorrhage in preterm infants. Lancet 1(8174):896–898. https://doi.org/10. 1016/s0140-6736(80)90836-3
- 12. Connor KM, SanGiovanni JP, Lofqvist C, Aderman CM, Chen J, Higuchi A et al (2007) Increased dietary intake of omega-3-polyunsaturated fatty acids reduces pathological retinal angiogenesis. Nat Med 13(7):868–873. https://doi.org/10.1038/nm1591
- Haggarty P (2010) Fatty acid supply to the human fetus. Annu Rev Nutr 30(1):237–255. https://doi.org/10.1146/annurev.nutr.012809.104742
- Khalesi N, Bordbar A, Khosravi N, Kabirian M, Karimi A (2018) The efficacy of omega-3 supplement on prevention of retinopathy of prematurity in premature infants: a randomized double-blinded controlled trial. Curr Pharm Des 24(17):1845–1848. https://doi.org/10.2174/138161282466618 0601094849
- Bernabe-García M, Villegas-Silva R, Villavicencio-Torres A, Calder PC, Rodríguez-Cruz M, Maldonado-Hernández J et al (2019) Enteral docosahexaenoic acid and retinopathy of prematurity: a randomized clinical trial. J Parenter Enteral Nutr 43(7):874–882. https://doi.org/10.1002/jpen. 1497
- Hellström A, Nilsson AK, Wackernagel D, Pivodic A, Vanpee M, Sjöbom U et al (2021) Effect of enteral lipid supplement on severe retinopathy of prematurity: a randomized clinical trial. JAMA Pediatr 175(4):359–367. https://doi.org/10.1001/jamapediatrics.2020.5653
- Lineham JD, Smith RM, Dahlenburg GW, King RA, Haslam RR, Stuart MC et al (1986) Circulating insulin-like growth factor I levels in newborn premature and full-term infants followed longitudinally. Early Hum Dev 13(1):37–46. https://doi.org/10.1016/0378-3782(86)90096-4
- Hellström A, Perruzzi C, Ju M, Engstrom E, Hard AL, Liu JL et al (2001) Low IGF-I suppresses VEGF-survival signaling in retinal endothelial cells: direct correlation with clinical retinopathy of prematurity. Proc Natl Acad Sci USA 98(10):5804–5808. https://doi.org/10.1073/pnas.101113998
- Hellgren G, Löfqvist C, Hård AL, Hansen-Pupp I, Gram M, Ley D et al (2016) Serum concentrations of vascular endothelial growth factor in relation to retinopathy of prematurity. Pediatr Res 79(1–1):70–75. https://doi.org/10. 1038/pr.2015.181
- Ley D, Hallberg B, Hansen-Pupp I, Dani C, Ramenghi LA, Marlow N et al (2019) rhIGF-1/rhIGFBP-3 in preterm infants: a phase 2 randomized controlled trial. J Pediatr 206:56–65.e8. https://doi.org/10.1016/j.jpeds.2018. 10.033
- Larnkjær A, Mølgaard C, Michaelsen KF (2012) Early nutrition impact on the insulin-like growth factor axis and later health consequences. Curr Opin Clin Nutr Metab Care 15(3):285–292. https://doi.org/10.1097/MCO. 0b013e328351c472
- Lee J, Dammann O (2012) Perinatal infection, inflammation, and retinopathy of prematurity. Semin Fetal Neonatal Med 17(1):26–29. https://doi. org/10.1016/j.siny.2011.08.007
- Husain SM, Sinha AK, Bunce C, Arora P, Lopez W, Mun KS et al (2013) Relationships between maternal ethnicity, gestational age, birth weight, weight gain, and severe retinopathy of prematurity. J Pediatr 163(1):67– 72. https://doi.org/10.1016/j.jpeds.2012.12.038
- Hellström A, Smith LE, Dammann O (2013) Retinopathy of prematurity. Lancet 382(9902):1445–1457. https://doi.org/10.1016/S0140-6736(13) 60178-6
- Sears JE, Hoppe G, Ebrahem Q, Anand-Apte B (2008) Prolyl hydroxylase inhibition during hyperoxia prevents oxygen-induced retinopathy. Proc Natl Acad Sci USA 105(50):19898–19903. https://doi.org/10.1073/pnas. 0805817105

- Shah PK, Narendran V, Kalpana N (2012) Aggressive posterior retinopathy of prematurity in large preterm babies in South India. Arch Dis Child Fetal Neonatal Ed 97(5):F371–375. https://doi.org/10.1136/fetalneona tal-2011-301121
- Rahimi N (2012) The ubiquitin-proteasome system meets angiogenesis. Mol Cancer Ther 11(3):538–548. https://doi.org/10.1158/1535-7163. MCT-11-0555
- Hoppe G, Yoon S, Gopalan B, Savage AR, Brown R, Case K, et al (2016) Comparative systems pharmacology of HIF stabilization in the prevention of retinopathy of prematurity. Proc Natl Acad Sci 113(18):E2516–25. Available from: https://pnas.org/doi/full/10.1073/pnas.1523005113. [cited 21 June 2023]
- Shibuya M (2011) Vascular endothelial growth factor (VEGF) and its receptor (VEGFR) signaling in angiogenesis: a crucial target for anti- and pro-angiogenic therapies. Genes Cancer 2(12):1097–1105. https://doi. org/10.1177/1947601911423031
- Smith LE, Shen W, Perruzzi C, Soker S, Kinose F, Xu X et al (1999) Regulation of vascular endothelial growth factor-dependent retinal neovascularization by insulin-like growth factor-1 receptor. Nat Med 5(12):1390–1395. https://doi.org/10.1038/70963
- Hellström A, Engström E, Hård AL, Albertsson-Wikland K, Carlsson B, Niklasson A et al (2003) Postnatal serum insulin-like growth factor I deficiency is associated with retinopathy of prematurity and other complications of premature birth. Pediatrics 112(5):1016–1020. https://doi.org/10.1542/ peds.112.5.1016
- Sato T, Kusaka S, Shimojo H, Fujikado T (2009) Vitreous levels of erythropoietin and vascular endothelial growth factor in eyes with retinopathy of prematurity. Ophthalmology 116(9):1599–1603. https://doi.org/10. 1016/j.ophtha.2008.12.023
- Palmer EA, Flynn JT, Hardy RJ, Phelps DL, Phillips CL, Schaffer DB et al (1991) Incidence and early course of retinopathy of prematurity. The cryotherapy for retinopathy of prematurity cooperative group. Ophthalmology 98(11):1628–40. https://doi.org/10.1016/s0161-6420(91)32074-8
- Austeng D, Källen KBM, Hellström A, Jakobsson PG, Johansson K, Tornqvist K et al (2011) Screening for retinopathy of prematurity in infants born before 27 weeks' gestation in Sweden. Arch Ophthalmol 129(2):167– 172. https://doi.org/10.1001/archophthalmol.2010.346
- Daniel E, Quinn GE, Hildebrand PL, Ells A, Hubbard GB, Capone A et al (2015) Validated system for centralized grading of retinopathy of prematurity: telemedicine approaches to evaluating acute-phase retinopathy of prematurity (e-ROP) study. JAMA Ophthalmol 133(6):675–682. https://doi. org/10.1001/jamaophthalmol.2015.0460
- Chiang MF, Quinn GE, Fielder AR, Ostmo SR, Paul Chan RV, Berrocal A et al (2021) International classification of retinopathy of prematurity, third edition. Ophthalmology 128(10):e51–68. https://doi.org/10.1016/j.ophtha. 2021.05.031
- Campbell JP, Chiang MF, Chen JS, Moshfeghi DM, Nudleman E, Ruambivoonsuk P et al (2022) Artificial intelligence for retinopathy of prematurity: validation of a vascular severity scale against international expert diagnosis. Ophthalmology 129(7):e69–76. https://doi.org/10.1016/j. ophtha.2022.02.008
- Zepeda-Romero LC, Martinez-Perez ME, Ruiz-Velasco S, Ramirez-Ortiz MA, Gutierrez-Padilla JA (2011) Temporary morphological changes in plus disease induced during contact digital imaging. Eye 25(10):1337–1340. https://doi.org/10.1038/eye.2011.170
- Royal College of Paediatrics and Child Health (2022) UK Screening of retinopathy of prematuirty guideline. Available from: https://www.rcpch. ac.uk/sites/default/files/2022-12/FC61116\_Retinopathy\_Guidelines\_14. 12.22.pdf [cited 28 June 2023]
- 40. The STOP-ROP Multicenter Study Group (2000) Supplemental therapeutic oxygen for prethreshold retinopathy of prematurity (STOP-ROP): a rand-omized, controlled trial. I: primary outcomes. Pediatrics 105(2):295–310. https://doi.org/10.1542/peds.105.2.295
- SUPPORT Study Group of the Eunice Kennedy Shriver NICHD Neonatal Research Network (2010) Target ranges of oxygen saturation in extremely preterm infants. N Engl J Med 362(21):1959–69. https://doi.org/10.1056/ NEJMoa0911781
- 42. The BOOST II United Kingdom, Australia, and New Zealand Collaborative Groups (2013) Oxygen saturation and outcomes in preterm infants. N Engl J Med 368(22):2094–104. https://doi.org/10.1056/NEJMoa1302298

- Schmidt B, Whyte RK, Asztalos EV, Moddemann D, Poets C, Rabi Y et al (2013) Effects of targeting higher vs lower arterial oxygen saturations on death or disability in extremely preterm infants: a randomized clinical trial. JAMA 309(20):2111. https://doi.org/10.1001/jama.2013.5555
- National Institute of Health and Care Excellence (2019) Specialist neonatal respiratory care for babies born preterm (NG124). Available from: www.nice.org.uk/guidance/ng124 [cited 28 June 2023]
- 45. Early treatment for Retinopathy of Prematurity Cooperative Group (2003) Revised indications for the treatment of retinopathy of prematurity: results of the early treatment for retinopathy of prematurity randomized trial. Arch Ophthalmol 121(12):1684. https://doi.org/10.1001/archopht. 121.12.1684
- Mutlu FM, Sarici SU (2013) Treatment of retinopathy of prematurity: a review of conventional and promising new therapeutic options. Int J Ophthalmol 6(2):228–236. https://doi.org/10.3980/j.issn.2222-3959.2013. 02.23
- 47. Papadopoulos N, Martin J, Ruan Q, Rafique A, Rosconi MP, Shi E et al (2012) Binding and neutralization of vascular endothelial growth factor (VEGF) and related ligands by VEGF Trap, ranibizumab and bevacizumab. Angiogenesis 15(2):171–185. https://doi.org/10.1007/s10456-011-9249-6
- Mahmood S, Roberts SA, Aslam TM, Parkes J, Barugh K, Bishop PN et al (2015) Routine versus as-needed bevacizumab with 12-weekly assessment intervals for neovascular age-related macular degeneration: 92-week results of the GMAN trial. Ophthalmology 122(7):1348–1355. https://doi.org/10.1016/j.ophtha.2015.03.017
- Scott IU, VanVeldhuisen PC, Ip MS, Blodi BA, Oden NL, Awh CC et al (2017) Effect of bevacizumab vs aflibercept on visual acuity among patients with macular edema due to central retinal vein occlusion: the SCORE2 randomized clinical trial. JAMA 317(20):2072. https://doi.org/10.1001/ jama.2017.4568
- Sangroongruangsri S, Ratanapakorn T, Wu O, Anothaisintawee T, Chaikledkaew U (2018) Comparative efficacy of bevacizumab, ranibizumab, and aflibercept for treatment of macular edema secondary to retinal vein occlusion: a systematic review and network meta-analysis. Expert Rev Clin Pharmacol 11(9):903–916. https://doi.org/10.1080/17512433.2018. 1507735
- Jhaveri CD, Glassman AR, Ferris FL, Liu D, Maguire MG, Allen JB et al (2022) Aflibercept monotherapy or bevacizumab first for diabetic macular edema. N Engl J Med 387(8):692–703. https://doi.org/10.1056/NEJMo a2204225
- Mintz-Hittner HA, Kennedy KA, Chuang AZ (2011) Efficacy of intravitreal bevacizumab for stage 3+ retinopathy of prematurity. N Engl J Med 364(7):603–615. https://doi.org/10.1056/NEJMoa1007374
- Moshfeghi DM, Berrocal AM (2011) Retinopathy of prematurity in the time of bevacizumab: incorporating the BEAT-ROP results into clinical practice. Ophthalmology 118(7):1227–1228. https://doi.org/10.1016/j. ophtha.2011.04.028
- Sankar MJ, Sankar J, Chandra P (2018) Anti-vascular endothelial growth factor (VEGF) drugs for treatment of retinopathy of prematurity. Cochrane Database Syst Rev 2018(1):CD009734 Available from: http://doi.wiley. com/10.1002/14651858.CD009734.pub3 Cochrane Neonatal Group, editor [cited 21 June 2023]
- Chan-Ling T, Gole GA, Quinn GE, Adamson SJ, Darlow BA (2018) Pathophysiology, screening and treatment of ROP: a multi-disciplinary perspective. Prog Retin Eye Res 62:77–119. https://doi.org/10.1016/j.preteyeres. 2017.09.002
- Sato T, Wada K, Arahori H, Kuno N, Imoto K, Iwahashi-Shima C et al (2012) Serum concentrations of bevacizumab (Avastin) and vascular endothelial growth factor in infants with retinopathy of prematurity. Am J Ophthalmol 153(2):327–333.e1. https://doi.org/10.1016/j.ajo.2011.07.005
- Miyake T, Sawada O, Kakinoki M, Sawada T, Kawamura H, Ogasawara K et al (2010) Pharmacokinetics of bevacizumab and its effect on vascular endothelial growth factor after intravitreal injection of bevacizumab in Macaque eyes. Investig Opthalmology Vis Sci 51(3):1606. https://doi.org/ 10.1167/iovs.09-4140
- Roohipoor R, Torabi H, Karkhaneh R, Riazi-Eafahani M (2019) Comparison of intravitreal bevacizumab injection and laser photocoagulation for type 1 zone II retinopathy of prematurity. J Curr Ophthalmol 31(1):61–65. https://doi.org/10.1016/j.joco.2018.10.008
- 59. Hillier RJ, Connor AJ, Shafiq AE (2018) Ultra-low-dose intravitreal bevacizumab for the treatment of retinopathy of prematurity: a case series.

Br J Ophthalmol 102(2):260–264. https://doi.org/10.1136/bjophthalm ol-2017-310408

- 60. Wallace DK, Kraker RT, Freedman SF, Crouch ER, Bhatt AR, Hartnett ME et al (2020) Short-term outcomes after very low-dose intravitreous bevacizumab for retinopathy of prematurity. JAMA Ophthalmol 138(6):698. https://doi.org/10.1001/jamaophthalmol.2020.0334
- Freedman SF, Hercinovic A, Wallace DK, Kraker RT, Li Z, Bhatt AR et al (2022) Low- and very low-dose bevacizumab for retinopathy of prematurity: reactivations, additional treatments, and 12-month outcomes. Ophthalmology 129(10):1120–1128. https://doi.org/10.1016/j.ophtha. 2022.05.019
- 62. NHS England Clinical Commissioning Policy (2023) Ranibizumab in retinopathy of prematurity (2201) [230401P]. Available from: https://www. england.nhs.uk/wp-content/uploads/2023/05/2201-ranibizumab-in-retin opathy-of-prematurity-policy.pdf [cited 9 Aug 2023]
- European Medicines Agency (2023) Lucentis: EPAR product information. Available from: https://www.ema.europa.eu/en/documents/produ ct-information/lucentis-epar-product-information\_en.pdf [cited 9 Aug 2023]
- 64. Stahl A, Lepore D, Fielder A, Fleck B, Reynolds JD, Chiang MF et al (2019) Ranibizumab versus laser therapy for the treatment of very low birthweight infants with retinopathy of prematurity (RAINBOW): an open-label randomised controlled trial. Lancet 394(10208):1551–1559. https://doi. org/10.1016/S0140-6736(19)31344-3
- Wu WC, Shih CP, Lien R, Wang NK, Chen YP, Chao AN et al (2017) Serum vascular endothelial growth factor after bevacizumab or ranibizumab treatment for retinopathy of prematurity. Retina 37(4):694–701. https:// doi.org/10.1097/IAE.00000000001209
- 66. Fidler M, Fleck BW, Stahl A, Marlow N, Chastain JE, Li J et al (2020) Ranibizumab population pharmacokinetics and free VEGF pharmacodynamics in preterm infants with retinopathy of prematurity in the RAINBOW trial. Transl Vis Sci Technol 9(8):43. https://doi.org/10.1167/tvst.9.8.43
- 67. Stahl A, Bründer M, Lagrèze WA, Molnár FE, Barth T, Eter N, et al (2022) Ranibizumab in retinopathy of prematurity – one-year follow-up of ophthalmic outcomes and two-year follow-up of neurodevelopmental outcomes from the CARE-ROP study. Acta Ophthalmol 100(1)e91-e99. Available from: https://onlinelibrary.wiley.com/doi/10.1111/aos.14852 [cited 21 June 2023]
- Sharma YR, Tripathy K, Venkatesh P, Gogia V (2014) Aflibercept how does it compare with other anti-VEGF drugs? Austin J Clin Ophthalmol 1(3):1016
- 69. Regeneron Pharmaceuticals, Inc (2023) EYLEA® (AFLIBERCEPT) Injection approved as the first pharmacologic treatment for preterm infants with Retinopathy of Prematurity (ROP) by the FDA. Available from: https:// investor.regeneron.com/news-releases/news-release-details/eylear-aflib ercept-injection-approved-first-pharmacologic [cited 9 Aug 2023]
- 70. European Medicines Agency (2023) Eylea: EPAR product information. Available from: https://www.ema.europa.eu/en/documents/productinformation/eylea-epar-product-information\_en.pdf
- Stahl A, Sukgen EA, Wu WC, Lepore D, Nakanishi H, Mazela J et al (2022) Effect of intravitreal aflibercept vs laser photocoagulation on treatment success of retinopathy of prematurity: the FIREFLEYE randomized clinical trial. JAMA 328(4):348. https://doi.org/10.1001/jama.2022.10564
- 72. Fleck BW, Reynolds JD, Zhu Q, Lepore D, Marlow N, Stahl A et al (2022) Time course of retinopathy of prematurity regression and reactivation after treatment with ranibizumab or laser in the RAINBOW trial. Ophthalmol Retina 6(7):628–637. https://doi.org/10.1016/j.oret.2022.02.006
- Snyder LL, Garcia-Gonzalez JM, Shapiro MJ, Blair MP (2016) Very late reactivation of retinopathy of prematurity after monotherapy with intravitreal bevacizumab. Ophthalmic Surg Lasers Imaging Retina 47(3):280–283. https://doi.org/10.3928/23258160-20160229-12
- 74. Chang E, Josan AS, Purohit R, Patel CK, Xue K (2022) A network metaanalysis of retreatment rates following bevacizumab, ranibizumab, aflibercept, and laser for retinopathy of prematurity. Ophthalmology 129(12):1389–1401. https://doi.org/10.1016/j.ophtha.2022.06.042
- Iwahashi C, Utamura S, Kuniyoshi K, Sugioka K, Konishi Y, Wada N et al (2021) Factors associated with reactivation after intravitreal bevacizumab or ranibizumab therapy in infants with retinopathy of prematurity. Retina 41(11):2261–2268. https://doi.org/10.1097/IAE.000000000003196
- Hamad AE, Moinuddin O, Blair MP, Schechet SA, Shapiro MJ, Quiram PA et al (2020) Late-onset retinal findings and complications in untreated

retinopathy of prematurity. Ophthalmol Retina 4(6):602–612. https://doi.org/10.1016/j.oret.2019.12.015

- Lepore D, Quinn GE, Molle F, Orazi L, Baldascino A, Ji MH et al (2018) Follow-up to age 4 years of treatment of type 1 retinopathy of prematurity intravitreal bevacizumab injection versus laser: fluorescein angiographic findings. Ophthalmology 125(2):218–226. https://doi.org/10.1016/j. ophtha.2017.08.005
- Hellström A, Hård AL, Engström E, Niklasson A, Andersson E, Smith L et al (2009) Early weight gain predicts retinopathy in preterm infants: new, simple, efficient approach to screening. Pediatrics 123(4):e638–645. https://doi.org/10.1542/peds.2008-2697
- Binenbaum G, Bell EF, Donohue P, Quinn G, Shaffer J, Tomlinson LA et al (2018) Development of modified screening criteria for retinopathy of prematurity: primary results from the postnatal growth and retinopathy of prematurity study. JAMA Ophthalmol 136(9):1034. https://doi.org/10. 1001/jamaophthalmol.2018.2753
- Moorthy S, Adams GGW, Smith G, Biswas S, Aliyan W, Bhudia R, et al (2023) Validation of the East London retinopathy of prematurity algorithm to detect treatment-warranted retinopathy of prematurity: a cohort study. Br J Ophthalmol bjo-2022-322522. https://doi.org/10.1136/ bjo-2022-322522
- Ramanathan A, Athikarisamy SE, Lam GC (2023) Artificial intelligence for the diagnosis of retinopathy of prematurity: a systematic review of current algorithms. Eye Lond Engl 37(12):2518–2526. https://doi.org/10. 1038/s41433-022-02366-y
- Filippi L, Cavallaro G, Berti E, Padrini L, Araimo G, Regiroli G et al (2019) Propranolol 0.2% eye micro-drops for retinopathy of prematurity: a prospective phase IIB study. Front Pediatr 7:180. https://doi.org/10.3389/ fped.2019.00180

# **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

# Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- ► Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at > springeropen.com